

# Quantum Mechanics is Linear Algebra

Noah Graham  
Middlebury College  
June 11, 2011

## Linear Algebra Cheat Sheet

Column vector (quantum state):  $|v\rangle = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \end{pmatrix}$

Row vector (dual state):  $\langle w| = (w_1^* \quad w_2^* \quad \dots)$

Inner product:  $\langle w|v\rangle = w_1^*v_1 + w_2^*v_2 + \dots$

Linear operator:  $\hat{A} = \begin{pmatrix} a_{11} & a_{12} & \dots \\ a_{21} & a_{22} & \\ \vdots & & \dots \end{pmatrix}$

Eigenvector & eigenvalue:  $\hat{A}|\lambda\rangle = \lambda|\lambda\rangle$


Average (expectation value):  $\langle A \rangle_\psi = \langle \psi | \hat{A} | \psi \rangle$

Standard deviation (uncertainty):  $\Delta A_\psi = \sqrt{\langle (\hat{A} - \langle A \rangle_\psi)^2 \rangle_\psi}$

operator	eigenstates	eigenvalues
$\hat{O}_{\uparrow} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$	$ \uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ $ \leftrightarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$	1 0
$\hat{O}_{\leftrightarrow} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$	$ \leftrightarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ $ \uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$	1 0
$\hat{O}_{\nearrow} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$	$ \nearrow\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ $ \searrow\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$	1 0
$\hat{O}_{\searrow} = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$	$ \searrow\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ $ \nearrow\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$	1 0
$\hat{O}_{\theta} = \begin{pmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{pmatrix}$ [the general case]	$ 1\rangle = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$ $ 0\rangle = \begin{pmatrix} \sin \theta \\ -\cos \theta \end{pmatrix}$	1 0

## The Physics

Electromagnetic waves (radio waves, microwaves, visible light, X-rays, gamma rays, etc.) have a **polarization**: a vector that lives in the two-dimensional vectorspace **perpendicular** to the direction of the wave's propagation. It gives the **axis** along which the wave's electric field oscillates.



[Image from Georgia State University HyperPhysics]

We'll hold fixed the frequency, wavelength and direction of propagation of the wave, so that we just focus on **polarization**.


## Working with polarization

A **polarizing filter** only allows light polarized along one axis to pass through. More precisely, it **projects** the polarization onto this axis — which only **reduces** the amount of light getting through.

Parallel: **all** gets through


Perpendicular: **none** gets through

(In between: **some** gets through)



[Image from Polymers and Liquid Crystals, Case Western Reserve University]

Note that the polarization vector can also be **complex**: Then the two components are out of phase and we get **circular** polarization.




[Image from Georgia State University HyperPhysics]

## A curious example

If vertically polarized light is incident on a horizontal polarizer, **nothing** gets through.

But if vertically polarized light is incident on a  $45^\circ$  polarizer and then a horizontal polarizer, some **does** get through.

Even though a polarizer **only blocks** light, it can cause **more** light to get through!



Note: The **intensity** of light (how much **energy** it carries) is proportional to the **square** of the length of these vectors.

**The problem:** In **quantum mechanics**, light comes in **discrete units**, known as **photons**. Each photon has to make its own decision: Do I get through or not?

## Postulate #1:

The **state of a quantum system** is completely described by a **normalized vector**  $|\psi\rangle$  in a (complex) vectorspace.

- The vectorspace may be finite or infinite dimensional.
- There are no “**hidden variables**” — there is no other information available about the system besides what we can find out from this vector.

---

Vertical polarization:  $|\uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$

Horizontal polarization:  $|\leftrightarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

Example

Diagonal polarization:  $|\nearrow\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

or  $|\searrow\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$

## Postulate #2:

Any physically measurable quantity is represented by a Hermitian linear operator  $\hat{O} = \hat{O}^\dagger$ .

- A Hermitian operator is one whose conjugate equals its transpose.
- Eigenvalues of a Hermitian operator are always real.
- Eigenvectors of a Hermitian operator form an orthonormal basis.

---

“Is polarization vertical?”:  $\hat{O}_\downarrow = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$

“Is polarization horizontal?”:  $\hat{O}_{\leftrightarrow} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$

“Is polarization diagonal?”:  $\hat{O}_{\nearrow} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

or  $\hat{O}_{\searrow} = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$

Example



### Postulate #3:

The possible results of a measurement of a physical quantity are the eigenvalues  $\lambda_i$  of the associated operator  $\hat{O}$ .

- Since  $\hat{O}$  is Hermitian, the eigenvalues are real.
- Physics is quantized!
- Many seemingly continuous physical quantities (energy, momentum, etc.) are actually the result of averaging discrete quantum outcomes over many particles.

---

$\hat{O}_{\uparrow}$ ,  $\hat{O}_{\leftrightarrow}$ ,  $\hat{O}_{\nearrow}$ , and  $\hat{O}_{\searrow}$  all have the same eigenvalues:

#### Example

- 1 = “yes” (photon gets through)
- 0 = “no” (photon blocked)

Note: No “maybe”!

## Postulate #4:

The **probability** of measuring the eigenvalue  $\lambda_i$  is  $P_i = |\langle \lambda_i | \psi \rangle|^2$ , where  $|\lambda_i\rangle$  is the **normalized eigenvector** of  $\hat{O}$  associated with the eigenvalue  $\lambda_i$ .

- $\langle \lambda_i | \psi \rangle$  is the  $i^{\text{th}}$  coordinate of  $|\psi\rangle$  in the basis  $\{|\lambda_i\rangle\}$ .
- Probabilities sum to the norm squared of  $|\psi\rangle$ , which is 1.
- If our state  $|\psi\rangle$  is **itself** an eigenstate  $|\lambda\rangle$  of  $\hat{O}$ , then we know for **certain** that the result of the measurement will be  $\lambda$ .

If  $|\psi\rangle$  is **not** an eigenstate, there are **multiple** possible outcomes.

### Example

- 
- Start in state  $|\uparrow\rangle$  and measure  $\hat{O}_{\downarrow}$ :  
Probability of “**yes**” is  $P_{\downarrow} = |\langle \downarrow | \uparrow \rangle|^2 = 1$  and probability of “**no**” is  $P_{\leftrightarrow} = |\langle \leftrightarrow | \uparrow \rangle|^2 = 0$ .
  - Start in state  $|\uparrow\rangle$  and measure  $\hat{O}_{\nearrow}$ :  
Probability of “**yes**” is  $P_{\nearrow} = |\langle \nearrow | \uparrow \rangle|^2 = \frac{1}{2}$  and probability of “**no**” is  $P_{\searrow} = |\langle \searrow | \uparrow \rangle|^2 = \frac{1}{2}$ .

## Postulate #5:

If we measure  $\hat{O}$  and find  $\lambda_i$ , then **immediately after that measurement**, the state **suddenly jumps** into the state  $|\lambda_i\rangle$ .

- Measurement **changes the system!**
- Two measurements in a row of the same quantity yield the same result.

- 
- If we start in state  $|\uparrow\rangle$  and measure  $\hat{O}_{\leftrightarrow}$ , we are **certain** to get the answer “no”:  $P_{\leftrightarrow} = 0$ .
  - But if we **first** measure  $\hat{O}_{\nearrow}$ , there is a 50% probability we get the answer “yes.” If so, we land in the state  $|\nearrow\rangle$ . Now a measurement of  $\hat{O}_{\leftrightarrow}$  has a 50% chance to yield the answer “yes” again. So

### Example

$$P_{\nearrow, \leftrightarrow} = |\langle \leftrightarrow | \nearrow \rangle|^2 \cdot |\langle \nearrow | \uparrow \rangle|^2 = \frac{1}{4} \neq 0$$

and now the photon **can** get through!

## Uncertainty

If we are in an **eigenstate** of the associated operator, we are **certain** of the outcome of a measurement of a physical quantity  $Q$  — it will be the associated eigenvalue.

If not, we can find the **average** of all the possible outcomes, weighted by their probabilities, and the associated **standard deviation**. Physicists call these the **expectation value**  $\langle Q \rangle_\psi$  and the **uncertainty**  $\Delta Q_\psi$  respectively.

But what if we would like to know about **two** physical quantities? If we would like to know both with **certainty**, we would need to be in an **eigenstate of both** operators at once.

**Theorem:** We can find a single basis of “simultaneous” eigenstates of  $\hat{A}$  and  $\hat{B}$  if and only if  $\hat{A}\hat{B} = \hat{B}\hat{A}$ .

Noncommutativity gives uncertainty!

## More on Uncertainty

The inner product obeys the **Schwarz inequality**

$$|\langle \phi | \psi \rangle|^2 \leq \langle \psi | \psi \rangle \langle \phi | \phi \rangle$$

For appropriate choices of  $|\phi\rangle$  and  $|\psi\rangle$ , a little algebra gives

$$\Delta A_\psi \cdot \Delta B_\psi \geq \frac{1}{2} |\langle \psi | (\hat{A}\hat{B} - \hat{B}\hat{A}) | \psi \rangle|$$

When the corresponding operators **don't commute**, there is a limit on how well we can know two physical quantities at once, because we have a clash of the **two different bases** of eigenstates we would need to work in to know each one with certainty.

Let's see how this works for the famous example. Now our vectorspace will consist of square-integrable ( $L^2$ ) functions. . .

## An Infinite-Dimensional Vectorspace

A particle moving on a line is specified by a **wavefunction**  $\psi(x)$ . Or by its Fourier transform,  $\tilde{\psi}(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{-ipx/\hbar} \psi(x) dx$ .

- $\psi(x)$  gives the state's coordinates in the **position** basis (specifies the function by its graph)
- $\tilde{\psi}(p)$  gives the state's coordinates in the **momentum** basis (specifies the function by its frequency spectrum)

Define the **inner product** on this vectorspace:

$$\langle \phi | \psi \rangle = \int_{-\infty}^{\infty} \phi(x)^* \psi(x) dx = \int_{-\infty}^{\infty} \tilde{\phi}(p)^* \tilde{\psi}(p) dp$$

Two **linear operators** on this vectorspace:

- $\hat{x} : \psi(x) \rightarrow x\psi(x)$  **or**  $\tilde{\psi}(p) \rightarrow i\hbar \frac{d}{dp} \tilde{\psi}(p)$ .
- $\hat{p} : \psi(x) \rightarrow -i\hbar \frac{d}{dx} \psi(x)$  **or**  $\tilde{\psi}(p) \rightarrow p\tilde{\psi}(p)$ .

## Infinite-Dimensional Uncertainty

The **commutator** of our two operators:

$$(\hat{x}\hat{p} - \hat{p}\hat{x}) : \psi(x) \rightarrow x \left( -i\hbar \frac{d}{dx} \psi(x) \right) - \left( -i\hbar \frac{d}{dx} (x\psi(x)) \right) = i\hbar \psi(x)$$

So  $(\hat{x}\hat{p} - \hat{p}\hat{x}) = i\hbar \cdot \hat{\mathbb{1}}$  and the uncertainty principle becomes

$$\Delta x \cdot \Delta p \geq \frac{\hbar}{2}$$

Just the mathematical statement that a **sharp spike** requires a **lot of frequencies**, or conversely a wave with a **narrow range of frequencies** must be **spread widely** throughout space.

This is why the sharp spikes in a high-speed data connection require a lot of **bandwidth** (many frequencies).

**Theorem:** It is only possible to have  $(\hat{x}\hat{p} - \hat{p}\hat{x}) = i\hbar \cdot \hat{\mathbb{1}}$  in an **infinite-dimensional** vector space.

Proof (one line) is left to the reader.

## Entangling Alliances

If we have **two** photons, our states are now  $|\uparrow\rangle|\uparrow\rangle$ ,  $|\uparrow\rangle|\leftrightarrow\rangle$ , etc.

Dimension of the space **multiplies** (tensor product).

Then we can have **entangled** states:

$$|\chi\rangle = \frac{1}{\sqrt{2}} (|\uparrow\rangle|\leftrightarrow\rangle + |\leftrightarrow\rangle|\uparrow\rangle)$$

Each photon has a 50% probability of being horizontal, a 50% probability of being vertical. But whatever we measure for the first photon, the second is **always opposite!**

Often applied to quantum **spin**, which works the same way with  $|\uparrow\rangle \Rightarrow |\uparrow\rangle$  (spin up) and  $|\leftrightarrow\rangle \Rightarrow |\downarrow\rangle$  (spin down). Note that **orthogonal** quantum states are now **180° apart** in the real world. (We have replaced  $SO(3)$  by  $SU(2)$ .)

Useful for quantum computation, quantum encryption, and more...



For more information:

<http://community.middlebury.edu/~ngraham/index.html#linearalgebra>